Author Index

Afonso, L.C.C., see Rumjanek, F.D., 145
Allsopp, B.A. and Allsopp, M.T.E.P.
Theileria parva: genomic DNA studies reveal intra-specific sequence diversity, 77
Allsopp, M.T.E.P., see Allsopp, B.A., 77
Amiri, P., Sakanari, J., Basch, P., Newport, G. and McKerrow, J.H.
The *Schistosomatium douthitti* cercarial elastase is biochemically and structurally distinct from that of *Schistosoma mansoni*, 113
Andrews, W.H., see Brothers, V.M., 235

Bachellerie, J.-P., see Gill, L.L., 69
Basch, P., see Amiri, P., 113
Beard, C.A., Wrightsman, R.A. and Manning, J.E.
Stage and strain specific expression of the tandemly repeated 90 kDa surface antigen gene family in *Trypanosoma cruzi*, 227
Beaumelle, B.D. and Vial, H.J.
Correlation of the efficiency of fatty acid derivatives in suppressing *Plasmodium falciparum* growth in culture with their inhibitory effect on acyl-CoA synthetase activity, 39
Berzins, K., see Snounou, G., 153
Bhaumik, D. and Datta, A.K.
Reaction kinetics and inhibition of adenosine kinase from *Leishmania donovani*, 181
Blum, J.J., see Darling, T.N., 121
Brindley, P.J., Gam, A.A., Pearce, E.J., Poindexter, R.W. and Neva, F.A.
Antigens from the surface and excretions/secretions of the filariform larva of *Strongyloides stercoralis*, 171
Brothers, V.M., Kuhn, I., Paul, L.S., Gabe, J.D., Andrews, W.H., Sias, S.R., McCaman, M.T., Dragon, E.A. and Files, J.G.
Characterization of a surface antigen of *Eimeria tenella* sporozoites and synthesis from a cloned cDNA in *Escherichia coli*, 235
Brown, K.N., see Snounou, G., 153
Bushell, G.R., Ingram, L.T., Fardoulys, C.A. and Cooper, J.A.
An antigenic complex in the rhoptries of *Plasmodium falciparum*, 105
Bushell, G.R., see Epping, R.J., 1

Campos, E.G., see Rumjanek, F.D., 145
Chakraborty, P. and Das, P.K.
Role of mannose/*N*-acetylglucosamine receptors in blood clearance and cellular attachment of *Leishmania donovani*, 55
Chakraborty, P.R., see Lindley, T.A., 135
Chappell, L., see Gill, L.L., 69
Clayton, C.E.
Most proteins, including fructose bisphosphate aldolase, are stable in the procyclic trypomastigote form of *Trypanosoma brucei*, 43
Cooper, J.A., see Bushell, G.R., 105
Cooper, J.A., see Epping, R.J., 1
Cross, G.A.M., see Gibbs, C.P., 197
Cummings, R.D., see Nyame, K., 265

Damian, R.T., see Nyame, K., 265
Darling, T.N. and Blum, J.J.
D-Lactate production by *Leishmania braziliensis* through the glyoxalase pathway, 121
Das, P.K., see Chakraborty, P., 55
Datta, A.K., see Bhaumik, D., 181
De la Cruz, V.F., Lal, A.A. and McCutchan, T.F.
Variation among circumsporozoite protein genes from rodent malarias, 31
De la Cruz, V.F., see McCutchan, T.F., 63
DeSimone, S.M., see Williams, S.A., 163
Dragon, E.A., see Brothers, V.M., 235

Edlind, T.D., see Lindley, T.A., 135
Elwood, H.J., see McCutchan, T.F., 63
Epping, R.J., Goldstone, S.D., Ingram, L.T., Upcroft, J.A., Ramasamy, R., Cooper, J.A., Bushell, G.R. and Geysen, H.M.
An epitope recognised by inhibitory monoclonal antibodies that react with a 51 kilodalton merozoite surface antigen in *Plasmodium falciparum*, 1

Fardoulys, C.A., see Bushell, G.R., 105
Files, J.G., see Brothers, V.M., 235
Fioravanti, C.F. and Kim, Y.
Rhodoquinone requirement of the *Hymenolepis diminuta* mitochondrial electron transport system, 129

Gabe, J.D., see Brothers, V.M., 235
Gam, A.A., see Brindley, P.J., 171
Geysen, H.M., see Epping, R.J., 1
Gibbs, C.P. and Cross, G.A.M.
Cloning and transcriptional analysis of a variant surface glycoprotein gene expression site in *Trypanosoma brucei*, 197
Gill, L.L., Hardman, N., Chappell, L., Qu, L.H., Nicoloso, M. and Bachellerie, J.-P.
Phylogeny of *Onchocera volvulus* and related species deduced from rRNA sequence comparisons, 69
Goldstone, S.D., see Epping, R.J., 1
Gunderson, J.H., see McCutchan, T.F., 63

Hardman, N., see Gill, L.L., 69
Hargreaves, J.J., see Howell, M.J., 21
Harn, D.A., see Oligino, L.D., 95
Hillman, Y., see Nicholls, S.C., 11
Holder, A.A., see Nicholls, S.C., 11
Holmquist, G., see Snounou, G., 153
Hori, E., see Nishina, M., 249

Howell, M.J. and Hargreaves, J.J.
Cloning and expression of *Taenia ovis* antigens in *Escherichia coli*, 21

Ingram, L.T., see Bushell, G.R., 105
Ingram, L.T., see Epping, R.J., 1

Kato, K., see Nishina, M., 249
Kennedy, M.W., see Knox, D.P., 207
Kim, Y., see Fioravanti, C.F., 129
King, D.L. and Turco, S.J.
A ricin agglutinin-resistant clone of *Leishmania donovani* deficient in lipophosphoglycan, 285
Knox, D.P. and Kennedy, M.W.
Proteinases released by the parasitic larval stages of *Ascaris suum*, and their inhibition by antibody, 207
Kuhn, I., see Brothers, V.M., 235

Lal, A.A., see De la Cruz, V.F., 31
Lal, A.A., see McCutchan, T.F., 63
Langsley, G., see Sharkey, A., 47
Lindley, T.A., Chakraborty, P.R. and Edlind, T.D.
Heat shock and stress response in *Giardia lamblia*, 135
Lockyer, M.J., see Nicholls, S.C., 11

Manning, J.E., see Beard, C.A., 227
Matsushita, K., see Nishina, M., 249
McCaman, M.T., see Brothers, V.M., 235
McCutchan, T.F., De la Cruz, V.F., Lal, A.A., Gunderson, J.H., Elwood, H.J. and Sogin, M.L.
Primary sequences of two small subunit ribosomal RNA genes from *Plasmodium falciparum*, 63
McCutchan, T.F., see De la Cruz, V.F., 31
McCutchan, T.F., see Vernick, K.D., 85
McKean, P.G., see Pritchard, D.I., 275
McKerrow, J.H., see Amiri, P., 113
McLean, A.P., see Sharkey, A., 47
McReynolds, L.A., see Williams, S.A., 163
Mercereau-Puijalon, O., see Sharkey, A., 47
Miller, R.L., Wang, A.L. and Wang, C.C.
Purification and characterization of the *Giardia lamblia* double-stranded RNA virus, 189

Neva, F.A., see Brindley, P.J., 171
Newport, G., see Amiri, P., 113
Nicholls, S.C., Hillman, Y., Lockyer, M.J., Odink, K.G. and Holder, A.A.
An S antigen gene from *Plasmodium falciparum* contains a novel repetitive sequence, 11
Nicoloso, M., see Gill, L.L., 69
Nishina, M., Hori, E., Matsushita, K., Takahashi, M., Kato, K. and Ohsaka, A.
^{1}H- and ^{13}C-NMR spectroscopic study of the metabolites in young adult *Angiostrongylus cantonensis* maintained in vitro, 249
Nyame, K., Cummings, R.D. and Damian, R.T.
Characterization of the high mannose asparagine-linked oligosaccharides synthesized by *Schistosoma mansoni* adult male worms, 265

Odink, K.G., see Nicholls, S.C., 11
Ohsaka, A., see Nishina, M., 249
Oligino, L.D., Percy, A.J. and Harn, D.A.
Purification and immunochemical characterization of a 22 kilodalton surface antigen from *Schistosoma mansoni*, 95

Patarapotikul, J., see Sharkey, A., 47
Paul, L.S., see Brothers, V.M., 235
Pearce, E.J., see Brindley, P.J., 171
Percy, A.J., see Oligino, L.D., 95
Pereira, M.E.A., see Prioli, R.P., 257
Perlmann, P., see Snounou, G., 153
Poindexter, R.W., see Brindley, P.J., 171
Prioli, R.P., Rosenberg, I., Shivakumar, S. and Pereira, M.E.A.
Specific binding of human plasma high density lipoprotein (cruzin) to *Trypanosoma cruzi*, 257
Pritchard, D.I., McKean, P.G. and Rogan, M.T.
Cuticle preparations from *Necator americanus* and their immunogenicity in the infected host, 275

Qu, L.H., see Gill, L.L., 69

Ramasamy, R., see Epping, R.J., 1
Roberts, S.M., see Saunders, N., 217
Rogan, M.T., see Pritchard, D.I., 275
Rosenberg, I., see Prioli, R.P., 257
Rumjanek, F.D., Campos, E.G. and Afonso, L.C.C.
Evidence for the occurrence of LDL receptors in extracts of schistosomula of *Schistosoma mansoni*, 145

Sakanari, J., see Amiri, P., 113
Saunders, N., Roberts, S.M. and Wilson, R.A.
Diazotised [^{125}I]iodosulphanilic acid is not a marker for the outer bilayer of the tegument of adult *Schistosoma mansoni*, 217
Sharkey, A., Langsley, G., Patarapotikul, J., Mercereau-Puijalon, O., McLean, A.P. and Walliker, D.
Chromosome size variation in the malaria parasite of rodents, *Plasmodium chabaudi*, 47
Shivakumar, S., see Prioli, R.P., 257
Sias, S.R., see Brothers, V.M., 235
Snounou, G., Viriyakosol, S., Holmquist, G., Berzins, K., Perlmann, P. and Brown, K.N.
Cloning of genomic fragment from *Plasmodium chabaudi* expressing a 105 kilodalton antigen epitope, 153
Sogin, M.L., see McCutchan, T.F., 63

Takahashi, M., see Nishina, M., 249
Turco, S.J., see King, D.L., 285

Upcroft, J.A., see Epping, R.J., 1

Vernick, K.D. and McCutchan, T.F.
Sequence and structure of a *Plasmodium falciparum* telomere, 85
Vial, H.J., see Beaumelle, B.D., 39
Viriyakosol, S., see Snounou, G., 153

Walliker, D., see Sharkey, A., 47
Wang, A.L., see Miller, R.L., 189
Wang, C.C., see Miller, R.L., 189
Williams, S.A., DeSimone, S.M. and McReynolds, L.A.
Species-specific oligonucleotide probes for the identification of human filarial parasites, 163
Wilson, R.A., see Saunders, N., 217
Wrightsman, R.A., see Beard, C.A., 227

Subject Index

Acyl-CoA synthetase, 39
Adenosine kinase, 181
Allergen, 171
Angiostrongylus cantonensis, 249
Antibody-mediated inhibition, 207
Antigen, 21, 171, 235
Antigen purification, 95
Antigenic complex, 105
Ascaris suum, 207

Binding, 145
Blood clearance, 55
Brugia malayi, 163
Brugia pahangi, 163

Cercaria, 113
Characterization of epitopes, 95
Chromosome polymorphism, 47
Circumsporozoite protein, 31
Collagen, 275
Cuticle, 275

Digitonin extraction, 217
DNA probe, 77, 163
cDNA library, 21, 235

Eimeria tenella, 235
Elastase, 113
Electron transport, 129
Enzyme kinetics, 181
Evolution, 31
Excretory/secretory product, 171
Expression, 21
Expression site associated gene, 197

Fatty acid, 39
Filariform larva, 171
Freeze-thaw, 217
Fructose bisphosphate aldolase, 43
Fumarate reductase, 129

Gene family, 227
Genetic polymorphism, 85
Genome size, 77
Giardia lamblia, 135, 189
D-[$^{13}C_6$]Glucose, 249
Glycoprotein, 55
Glyoxalase, 121
Growth inhibition, 39

Heat shock, 135
High density lipoprotein, 257
High mannose N-linked oligosaccharides, 265
Hymenolepis diminuta, 129

Immunogenicity, 275
In vitro-released products, 207
Infection, 275
Infective larvae (L2), 207
Inhibitor, 207
Irradiated cercariae, 95

D-Lactate, 121
Leishmania, 121
Leishmania donovani, 55, 181, 285
Lipophosphoglycan, 285
Lipoprotein, low density and very low density, 145
^{125}I-Lipoprotein, 145
Liposome, 145
Lung-stage larvae (L3/4), 207
Lymphatic filariasis, 163

Macrophage, 55, 285
Malaria, 1, 31, 47, 85, 153
Merozoite surface, 1
Metabolic end products, 249
Methylglyoxal, 121
Mitochondria, 129
Monoclonal antibody, 1, 105

NADH oxidase, 129
Necator americanus, 275
Nematode, 207
Neoglycoprotein, 55
Neuraminidase, 257
Nitrocellulose filter, 145
NMR spectroscopy, 249
Nucleoside analog phosphorylation, 181

Onchocerca volvulus, 69
Oligosaccharide synthesis, 265

Pc 105 antigen, 153
Pf 155 antigen, 153
pH optima, 207
Phylogeny, 69
Plasmid vector, 21
Plasmodium berghei, 31
Plasmodium chabaudi, 47, 153
Plasmodium falciparum, 1, 11, 39, 63, 85, 105, 153
Plasmodium yoelii, 31
Probe, 63
Protease, 113
Protein synthesis, 135
Proteinases, 207
Pulsed field gradient gel electrophoresis, 47

Radiolabelling, 217
Receptor, 145
Recombinant DNA, 153

Repeat region, 11
Repeated DNA, 163
Rhodoquinone, 129
Rhoptry, 105
Ribosomal RNA, 63
Ricin, 285
dsRNA virus, 189
rRNA sequence, 69

S antigen gene, 11
Saturation, 145
Schistosoma mansoni, 95, 265
Schistosoma douthitti, 113
Schistosome, 113
Schistosomula, 145
Sheep, 21
Species identification, 163
Sporozoite, 235
Stability, 43
Strain characterization, 77
Stress response, 135
Strongyloides stercoralis, 171
Substrate specificity, 207
Sugar receptor, 55
Surface antigen, 227
Surface membrane, 95
Swimmer's itch, 113

Taenia ovis, 21
Telomere, 85
Theileria parva, 77
Triton X-100, 145
Trypanosoma brucei, 43, 197
Trypanosoma cruzi, 227, 257

Variant surface glycoprotein expression, 197
Virus, 189

Worm surface, 171